

THE ENTROPY OF LEAD

Computational Document

by

George Hardgrove
Chemistry Department
St. Olaf College
Northfield, MN 55057

hardgrov@lars.acc.stolaf.edu

The instructions for using this document are in the file cppbins.mcd.

© Copyright George Hardgrove 1996-2004. All rights reserved. You are welcome to use this document in your own classes but commercial use is not allowed without the permission of author. Mathcad 2001i version by Theresa Julia Zielinski, Monmouth University

The heat capacity data are located in a file called cplead. Values of Cp and absolute temperature are read into the computer. The computer then calculates values of $ds = Cp/T$ for each temperature.

```
A := READPRN("CPLEAD.prn" ndata := rows(A)    i := 0..ndata - 1
```

$T_i := (A^{(0)})_i$ $Cp_i := (A^{(1)})_i$ $dS_i := \frac{Cp_i}{T_i}$ This assigns the columns of the A matrix to temperature and heat capacity vectors.

Next view the contents of these vectors.

T =



	0
0	10
1	15
2	20
3	25
4	30
5	50
6	70
7	100
8	150
9	200
10	250
11	298

Cp =

	0
0	2.8
1	7
2	10.8
3	14.1
4	16.5
5	21.8
6	23.3
7	24.5
8	25.3
9	25.8
10	26.2
11	26.6

dS =

	0
0	0.28
1	0.4667
2	0.54
3	0.564
4	0.55
5	0.436
6	0.3329
7	0.245
8	0.1687
9	0.129
10	0.1048
11	0.0893



To use the simple formula for trapezoid integration with evenly spaced points, we need to use interpolation methods. First we split up the space between the first and last points into N segments where we can adjust N.

$$\Delta x := \frac{T_{\text{ndata}-1} - T_0}{N} \quad j := 0..N \quad \Delta x = 28.8 \quad x_j := T_0 + j \cdot \Delta x$$

The cspline function accumulates derivatives of the function at each point and puts them into a vector called vs.


$$vs := \text{cspline}(T, dS)$$

We can now find the value of the function at each x_j . On the graph below the black + signs are the experimental data, the red bars indicate the value of the function at evenly spaced points fx_j .

$$fx_j := \text{interp}(vs, T, dS, x_j) \quad zx_j := 0 \quad \text{zx}_j \text{ sets to lower level for the error bars.}$$

$$\text{Area} := \sum_{j=1}^{N-1} \Delta x \cdot fx_j + \left[\frac{(fx_0 + fx_N)}{2} \cdot \Delta x \right] \quad \text{The evenly spaced points are put into the area formula}$$

$$N \equiv 10 \quad \text{Area} = 60.9846$$



Now we need to consider the contribution to the entropy for the region from 0° K up to the lowest temperature at which the heat capacity was measured T_0 . In this region the heat capacity follows the function

$$C_p := a \cdot T^3 \quad \frac{C_p}{T} := a \cdot T^2 \quad \int_0^{T_0} \frac{C_p}{T} dT := \frac{C_p(T_0)}{3}$$

$$C_{p0} = 2.8$$

$$\Delta S := \frac{C_{p0}}{3} \quad \Delta S = 0.9333$$

Therefore the total entropy for 0o K to 298o K is Area + ΔS

$$\text{Area} + \Delta S = 61.9179$$

The literature value is 64.81 J/Kmol. Differences between the calculated and literature values occur because of the interpolation technique used in calculations.